



**Demo Report**

**Report Issued** 06/04/2013 12:19  
**Last Report Issued** 01/30/2013 14:51

**88888**

Clinical Example Report  
Attn: IT Department  
200 Welsh Road  
Horsham, PA 19044-2208

**Patient Name** 1900SP  
**Patient ID** 1900SP  
**Chain** 13000600  
**Age** Not Given **Gender** Not Given

**Workorder** 13000600  
**Received** 01/30/2013 12:30

**Sample ID** 13000600-001  
**Matrix** Serum or Plasma  
**Patient Name** 1900SP  
**Patient ID** 1900SP  
**Container Type** Clear vial

**Collect Dt/Tm** Not Given  
**Source** Not Given

**Approx Vol/Weight** Not Given

**Receipt Notes** None Entered

Analysis and Comments	Result	Units	Reporting Limit	Notes
-----------------------	--------	-------	-----------------	-------

**1900SP Dyazide, Serum/Plasma**

Analysis by High Performance Liquid Chromatography/Tandem Mass Spectrometry (LC-MS/MS)

Triamterene Synonym(s): Dyrenium® Chronic oral administration of 75 mg triamterene in combination with 50 mg hydrochlorothiazide resulted in peak plasma concentrations of 200 +/- 130 ng/mL triamterene.	None Detected	ng/mL	5.0	
---	---------------	-------	-----	--

Hydrochlorothiazide Synonym(s): Microzide® Daily administration of oral hydrochlorothiazide produced the following plasma concentrations in hypertensive patients:  25 mg = 17 +/- 8 ng/mL trough and 76 +/- 26 ng/mL 5 hours post dose  75 mg = 34 +/- 17 ng/mL trough and 200 +/- 93 ng/mL 5 hours post dose.	None Detected	ng/mL	25	
---	---------------	-------	----	--